

This product is for research use only (not for diagnostic or therapeutic use)

contact: support@agrisera.com

Agrisera AB | Box 57 | SE-91121 Vännäs | Sweden | +46 (0)935 33 000 | www.agrisera.com

Product no AS13 2754

CDPK | Calcium-dependent protein kinase isoforms (1,2,3,4,7,8,10,11,14,20,25,30,32)

Product information

Immunogen KLH-conjugated synthetic peptide conserved in isoform 1,2,3,4,7,8,10,11,14,20,25,30,32 of Arabidopsis thaliana CDPK

Host Rabbit

Clonality Polyclonal

Purity Immunogen affinity purified serum in PBS pH 7.4.

Format Lyophilized

Quantity 50 μg

Reconstitution For reconstitution add 50 μl of sterile water

Storage Store lyophilized/reconstituted at -20°C; once reconstituted make aliquots to avoid repeated freeze-thaw cycles. Please remember to spin the tubes briefly prior to opening them to avoid any losses that might occur from material adhering to

the cap or sides of the tube.

Application information

Recommended dilution 1:1000 (WB)

Expected | apparent MW 52-55 kDa

Confirmed reactivity Arabidopsis thaliana, Hordeum vulgare, Populus sp., Triticum aestivum

Predicted reactivity Fragaria ananassa, Malus domestica, Nicotiana attenuata, Oryza sativa, Vicia faba

Species of your interest not listed? Contact us

Not reactive in No confirmed exceptions from predicted reactivity are currently known

Selected references Kumar at al. (2024). Characterization of putative calcium-dependent protein kinase-1 (TaCPK-1) gene: hubs in

signalling and tolerance network of wheat under terminal heat.Biotech, Volume 14, article number 150.

 $\underline{Siddigui} \ et \ al. \ (2020). \ Melatonin \ and \ calcium \ function \ synergistically \ to \ promote \ the \ resilience \ through \ ROS \ metabolism$

under arsenic-induced stress. Journal of Hazardous MaterialsVolume 398, 5 November 2020, 122882

Ciesla et al. (2016) A Role for Barley Calcium-Dependent Protein Kinase CPK2a in the Response to Drought. Front

Plant Sci. 2016 Oct 25;7:1550.