

This product is for research use only (not for diagnostic or therapeutic use)

contact: support@agrisera.com

Agrisera AB | Box 57 | SE-91121 Vännäs | Sweden | +46 (0)935 33 000 | www.agrisera.com

## Product no AS09 604

## Donkey anti-Sheep IgG (H&L), HRP conjugated

## **Product information**

**Immunogen** Purified sheep IgG, whole molecule

**Host** Donkey

Clonality Polyclonal

**Purity** Immunogen affinity purified donkey IgG.

Format Lyophilized

Quantity 1 mg

Reconstitution For reconstitution add 1,1 ml of sterile water, Let it stand 30 minutes at room temperature to dissolve, Centrifuge to

remove any particulates, Prepare fresh working dilutions daily

Storage Store lyophilized material at 2-8 °C. For storage at -20 °C after reconstitution dilute antibody solution with an equal volume of glycerol to obtain final glycerol concentration of 50 % to prevent loss of enzymatic activity. Such solution will not freeze in -20°C. If you are using a 1:5000 dilution prior to diluting with glycerol, then you would need to use a

1:2500 dilution after adding glycerol. Prepare working dilution prior to use and then discard, Be sure to mix well but

**Additional information** 

Antibody concentration is 1 mg/ml and it has been purified by antigen-specific chromatography

HRP-conjugate is supplied in 10 mM Sodium Phosphate, 0.15 M Sodium Chloride, pH 7.2, 10 % (w/v) BSA, Protease/IgG free

0.1 % (v/v) of ProClin 150 is used as preservative.

For blocking: BSA and nonfat milk have to be replaced by other blocking reagents, like doneky serum or commercial formulations which are free from bovine IgG.

## **Application information**

Recommended dilution 1:200 -1:5 000 (ELISA), 1:200-1:5 000 (WB) 1:500 -1:5000 (IHC)

Confirmed reactivity Sheep IgG heavy chains and light chains on all sheep immunoglobulins

Not reactive in non-immunoglobulin sheep serum proteins based in immunoelectrophoresis