


Product no **AS19 4252****eIF5 | Eukaryotic initiation factor eIF5****Product information**

<b>Immunogen</b>	Recombinant eIF5 of <i>Triticum aestivum</i>
<b>Host</b>	Rabbit
<b>Clonality</b>	Polyclonal
<b>Purity</b>	Serum
<b>Format</b>	Lyophilized
<b>Quantity</b>	50 µl
<b>Reconstitution</b>	For reconstitution add 200 µl of sterile water
<b>Storage</b>	Store lyophilized/reconstituted at -20°C; once reconstituted make aliquots to avoid repeated freeze-thaw cycles. Please remember to spin the tubes briefly prior to opening them to avoid any losses that might occur from material adhering to the cap or sides of the tube.

**Application information**

<b>Recommended dilution</b>	1 : 8000 (WB)
<b>Expected   apparent MW</b>	49   50 kDa
<b>Confirmed reactivity</b>	<i>Arabidopsis thaliana</i> , <i>Lactuca sativa</i> , <i>Nicotiana tabacum</i> , <i>Triticum aestivum</i> , <i>Pisum sativum</i>
<b>Predicted reactivity</b>	<i>Nelumbo nucifera</i> , <i>Triticum urartu</i> , <i>Zea mays</i> Species of your interest not listed? <a href="#">Contact us</a>

**Application example**

Recommend incubation for 1 hr at room temperature in 5% non-fat dry milk in phosphate-buffer saline with 0.5% Tween-80 (PBST). Figure shows plant leaf extracts or wheat germ extract (5 µl of the supernatant of 50 mg liquid nitrogen powdered tissue/100 µl 2X SDS Laemli buffer containing protease inhibitor crushed with a plastic pestle in a microfuge tube and centrifuged for 10 min at 4 C.) Image shown is using Agrisera Anti-rabbit HRP (1/30,000) for 1 hr at RT in 5% non-fat dry milk in phosphate-buffer saline with 0.5% Tween-80 (PBST) and using Agrisera Ultra Bright substrate and exposed for 5 seconds in a BioRad ChemDoc on high resolution. Blue star indicates eIF5 band. Courtesy Dr. Karen Browning, The University of Texas at Austin, USA