

**Product no** **AS16 3339****Goat anti-Human IgG Fc, ALP conjugated, min, cross-reactivity to human IgA + IgM, bovine, horse, mouse or rabbit serum proteins****Product information****Immunogen** | Purified Human IgG, Fc fragment**Host** | Goat**Clonality** | Polyclonal**Purity** | Immunogen affinity purified goat IgG.**Format** | Liquid**Quantity** | 1,0 mg**Storage** | Store non-diluted antibody at 2-8°C. For storage at -20 °C, dilute with an equal volume of glycerol to prevent loss of enzymatic activity. Prepare working dilutions prior to use and then discard.**Additional information** | Antibody purity is > 95% based on SDS-PAGE.  
Affinity purified using solid phase Human IgG .  
Antibody is supplied in 30 mM Triethanolamine, pH 7.2, 5 mM Magnesium Chloride, 0.1 mM Zinc Chloride, 1 % (w/v) BSA, Protease/IgG free. 0.05% (w/v) Sodium Azide is added as a preservative.

Based on IEP, this antibody reacts with: • heavy ( ) chains on human IgG .

Based on IEP, no reactivity is observed to: • non-immunoglobulin human serum immunoglobulins • light chains on all human immunoglobulins • human IgA or IgM • serum proteins from bovine, mouse, or rabbit .

**Application information****Recommended dilution** | 1 : 20-1 : 2000 (Flow cyt), (IL)**Confirmed reactivity** | Human IgG Fc