

This product is for research use only (not for diagnostic or therapeutic use)

contact: support@agrisera.com

Agrisera AB | Box 57 | SE-91121 Vännäs | Sweden | +46 (0)935 33 000 | www.agrisera.com

Product no AS12 2105

SRP43 | Signal recognition peptide 43

Product information

Immunogen KLH-conjugated synthetic peptide derived from Arabidopsis thaliana SRP43, UniProt: O22265, TAIR: AT2G47450

Host Chicken

Clonality Polyclonal

Purity Immunogen affinity purified serum in PBS pH 7.4.

Format Lyophilized

Quantity 50 μg

Reconstitution For reconstitution add 50 μl of sterile water

Storage Store lyophilized/reconstituted at -20°C; once reconstituted make aliquots to avoid repeated freeze-thaw cycles. Please, remember to spin tubes briefly prior to opening them to avoid any losses that might occur from lyophilized

material adhering to the cap or sides of the tubes.

Application information

Recommended dilution 1:5000 (WB)

Expected | apparent 41.3

41.3 kDa

Not reactive in no confirmed exceptions from predicted reactivity are currently known

Selected references To be added when available, antibody released in June 2023.



15 μ g/well of total protein extracted freshly from rosetta leaves of *Arabidopsis thaliana* with 2X Laemmli buffer and denatured at 95 °C/5 min. Samples were separated in the cold on 12% SDS-PAGE and blotted for 1.5 h to nitrocellulose (pore size of 0.45 μ m), using: wet, semi-dry transfer in the cold. Blot was blocked with 5 % milk for: 1h/RT with agitation. Blot was incubated in the primary antibody at a dilution of 1: 1000 in TBS for ON/4°C with agitation. The antibody solution was decanted, and the blot was rinsed briefly twice, then washed once for 5 min in TBS-T (1% Tween-20) and 2 times for 10 min in TBS at RT with agitation. Blot was incubated in matching secondary antibody (anti-chicken IgG horse radish peroxidase conjugated AS10 1489, Agriera) diluted to 1: 5000 in for 1.5 h/RT with agitation. The blot was washed as above and developed with a following chemiluminescent detection reagent: <u>AS16 ECL-N-10</u> AgriseraBright (Agrisera). Exposure time was 1 minute.

Courtesy od dr. Peng Wang, School of Biological Sciences, The University of Hong Kong