


Product no **AS14 2829****UAGPase | UDP-GlcNAc pyrophosphorylase****Product information**

<b>Immunogen</b>	His-tagged full length recombinant UAGPase of <i>Arabidopsis thaliana</i> , overexpressed and purified from <i>E.coli</i> , UniProt: <a href="#">O64765</a> , TAIR: <a href="#">AT2G35020</a> . Sequence used for immunization is conserved in both isoforms: UDP-N-acetylglucosamine diphosphorylase 1 ( <a href="#">Q940S3</a> ) and ( <a href="#">O64765</a> )
<b>Host</b>	Rabbit
<b>Clonality</b>	Polyclonal
<b>Purity</b>	Serum
<b>Format</b>	Lyophilized
<b>Quantity</b>	50 µl
<b>Reconstitution</b>	For reconstitution add 50 µl of sterile water
<b>Storage</b>	Store lyophilized/reconstituted at -20°C; once reconstituted make aliquots to avoid repeated freeze-thaw cycles. Please remember to spin the tubes briefly prior to opening them to avoid any losses that might occur from material adhering to the cap or sides of the tube.

**Application information**

<b>Recommended dilution</b>	1 : 10 000 (WB)
<b>Expected   apparent MW</b>	55,76 kDa
<b>Confirmed reactivity</b>	<i>Arabidopsis thaliana</i> , <i>Hordeum vulgare</i> , <i>Nicotiana tabacum</i>
<b>Predicted reactivity</b>	<i>Glycine soja</i> , <i>Medicago truncatula</i> , <i>Morus notabilis</i> , <i>Populus trichocarpa</i> , <i>Theobroma cacao</i> , <i>Zea mays</i> , for more species, please inquire Species of your interest not listed? <a href="#">Contact us</a>
<b>Not reactive in</b>	No confirmed exceptions from predicted reactivity are currently known
<b>Additional information</b>	This antibody is recognizing recombinant UAGPase at 0,25 pmol
<b>Selected references</b>	<a href="#">Fernández-San Millán et al. (2018)</a> . Physiological Performance of Transplastomic Tobacco Plants Overexpressing Aquaporin AQP1 into Chloroplast Membranes. <i>J Exp. Bot.</i> ery148, <a href="https://doi.org/10.1093/jxb/ery148">https://doi.org/10.1093/jxb/ery148</a> . <a href="#">Kleczkowski LA &amp; Decker DD (2015)</a> Sugar activation for production of nucleotide sugars as substrates for glycosyltransferases in plants. <i>J. Appl. Glycosci.</i> 2015 Volume 62 Issue 2 Pages 25-36.

**Application example**

10 µg of total protein from *Arabidopsis thaliana* leaf (1) *Hordeum vulgare* leaf (2), *Nicotiana tabacum* (3), and recombinant UAGPase 0.25 pmol (R), were extracted with Protein Extraction Buffer PEB ([AS08 300](#)). Samples were diluted with 1X sample buffer (NuPAGE LDS sample buffer (Invitrogen) supplemented with 50 mM DTT and heat at 70°C for 5 min and kept on ice before loading. Protein samples were separated on 4-12% Bolt Plus gels, LDS-PAGE and blotted for 70 minutes to PVDF using tank transfer. Blots were blocked immediately following transfer in 2% blocking reagent or 5% non-fat milk dissolved in 20 mM Tris, 137 mM sodium chloride pH 7.6 with 0.1% (v/v) Tween-20 (TBS-T) for 1h at room temperature with agitation. Blots were incubated in the primary antibody at a dilution of 1: 10 000 (in blocking reagent) for 1h at room temperature with agitation. The antibody solution was decanted and the blot was rinsed briefly twice, and then washed 1x15 min and 3x5 min with TBS-T at room temperature with agitation. Blots were incubated in secondary antibody (anti-rabbit IgG horse radish peroxidase conjugated, recommended secondary antibody [AS09 602](#), Agrisera) diluted to 1:25 000 in blocking reagent for 1h at room temperature with agitation. The blots were washed as above. The blot was developed for 5 min with detection reagent according to the manufacturers instructions. Images of the blots were obtained using a CCD imager (VersaDoc MP 4000) and Quantity One software (Bio-Rad). Exposure time was 10 seconds for



This product is **for research use only** (not for diagnostic or therapeutic use)

**contact: [support@agrisera.com](mailto:support@agrisera.com)**

Agrisera AB | Box 57 | SE-91121 Vännäs | Sweden | +46 (0)935 33 000 | [www.agrisera.com](http://www.agrisera.com)

recombinant UAGpase and 1 minute for plant extracts.